## **Towards Three-Dimensional Imaging of Yeast Spores Using Coherent X-ray Diffraction Microscopy**

*Adrian P. Mancuso*<sup>1</sup>, Changyong Song<sup>1</sup>, Huaidong Jiang<sup>1</sup>, Damien Ramunno-Johnson<sup>1</sup>, Tetsuya Ishikawa<sup>2</sup>, Fuyu Tamanoi<sup>3</sup> and Jianwei Miao<sup>1</sup>

The three-dimensional imaging of whole cells at around 10 nm resolution is of great biological interest. Using coherent x-ray diffraction microscopy, we demonstrate high-resolution imaging of an unlabelled yeast spore at different angles of tilt. The experiments were carried out on an undulator beamline at SPring-8, Japan. The collected diffraction data were analyzed using an explicit account for detector pixel size. Reconstructions of projections of the yeast spore, at differing angles, have been performed based on a guided hybrid input-output algorithm (GHIO). These reconstructions are shown, along with an account of the reconstruction method. This work represents a practical step toward routine three-dimensional imaging of unlabelled biological cells using X-ray diffraction microscopy.

This work was supported by the National Science Foundation (NSF), the Department of Energy (DOE), and the California NanoSystems Institute (CNSI). Use of the RIKEN beamline, BL29XUL at SPring-8 was supported by RIKEN.

<sup>&</sup>lt;sup>1</sup>Department of Physics and Astronomy, University of California, Los Angeles, CA 90095 USA

<sup>&</sup>lt;sup>2</sup>SPring-8/RIKEN, 1-1-1, Kouto, Mikazuki, Sayo-gun, Hyogo 679-5198, Japan.

<sup>&</sup>lt;sup>3</sup>Department of Microbiology, Immunology and Molecular Genetics, Jonsson Comprehensive Cancer Center, Molecular Biology Institute, University of California, Los Angeles, CA 90095 USA